



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

is true of  $x_2$  and  $x_2'$ . Hence,  $\infty$  is a root of the sextic covariant, whose leading coefficient must therefore vanish. This gives  $8a_0^2a_3 - 4a_0a_1a_2 + a_1^3 = 0$  as the required condition. The same result may be readily obtained by elementary methods.

Also solved by HERBERT N. CARLETON, OTTO J. RAMLER, and HORACE OLSON.

E. B. ESCOTT sent in three solutions of 515 (Geometry). The first two solutions were solutions given in E. CATALAN'S *Théorèmes et Problèmes de Géométrie Élémentaire*, pp. 237-239. The third is a neat original solution.

**516 (Geometry).** Proposed by R. M. MATHEWS, Riverside, California.

Through the edges of a trihedral angle planes are passed orthogonal to the opposite faces. Prove them coaxial.

#### I. SOLUTION BY NATHAN ALTSHILLER, University of Oklahoma.

The planes  $x, y, z$  which pass through the edges  $OA, OB, OC$  of the trihedral angle  $O-ABC$  and which are perpendicular to the opposite faces  $a, b, c$ , cut these faces along lines  $OD, OE, OF$ , respectively. A plane  $p$  perpendicular to  $OA$  cuts the edges  $OA, OB, OC$  in the points  $A, B, C$ , and the lines  $OD, OE, OF$  in the points  $D, E, F$ . (The reader may readily construct the figure.)

The plane  $x$  is perpendicular to  $a$  by construction, and to the plane  $p$ , because  $x$  passes through  $OA$ ; hence,  $x$  is perpendicular to the line of intersection  $BC$  of  $p$  with  $a$ , and therefore  $BC$  is perpendicular to  $AD$ .

The plane  $b$  is perpendicular to  $y$  by construction and to the plane  $p$ , because  $b$  passes through  $OA$ ; hence,  $b$  is perpendicular to the line of intersection  $BE$  of  $y$  with  $p$ , and, therefore,  $BE$  is perpendicular to  $AC$ . For similar reasons,  $CF$  is perpendicular to  $AB$ .

The three altitudes  $AD, BE, CF$  of the triangle  $ABC$  concur, according to a well-known proposition, in a point  $H$ , the orthocenter of  $ABC$ ; hence  $H$  is a common point of the planes  $x, y, z$ . Now these three planes have obviously the point  $O$  in common, hence they pass through the line  $OH$ .

Incidentally we have also proved: *The locus of the orthocenter of the triangle determined by three concurrent lines and a variable plane perpendicular to one of the given lines is a straight line concurrent with the given lines.*

#### II. SOLUTION BY W. WOOLSEY JOHNSON, Annapolis, Md.

Referred to the sphere, the problem is that of the existence of the orthocenter of the spherical triangle  $ABC$ .

Let  $CD = p$  be the perpendicular from  $C$  upon  $AB$  and let the perpendicular from  $A$  upon  $BC$  cut it in  $O$ .

From the right triangle  $AOD$  we have

$$\tan OD = \tan OAD \sin AD. \quad (1)$$

From  $AEB$ , we have

$$\cot OAD = \cos c \tan B. \quad (2)$$

From  $ADC$ , we have

$$\sin AD = \tan p \cot A. \quad (3)$$

Dividing (3) by (2) and substituting in (1), we have  $\tan OD = \tan p \cot A \cot B \sec c$ .

Interchanging  $A$  and  $B$ ,  $c$  and  $p$  remain unchanged; hence, the perpendicular from  $B$  cuts off from  $p$  the same segment  $OD$  or the three perpendiculars meet in a common point.

Also solved by HORACE OLSON, L. E. LUNN, C. J. PAYNE, WILLIAM HOOVER, and the PROPOSER.